arXiv:1303.3790v2 [physics.gen-ph] 13 May 2013

What the Casimir-effect really is telling about zero-point energy

Gerold Gründler*
Astrophysical Institute Neunhof - Obere Dorfstraße 18, D-90427 Nürnberg, Germany

The attractive force between metallic surfaces, predicted by Casimir in 1948, seems to indicate the physical existence and measurability of the quantized electromagnetic field's zero-point energy. It is shown in this article, that Casimir's derivation depends essentially on a misleading idealization. When that idealization is replaced by a realistic assumption, Casimir's argument turns to the exact opposite: The observed Casimir force does positively prove, that the electromagnetic field's zero-point energy does *not* exert forces onto metallic surfaces.

PACS numbers: 03.70.+k, 04.20.Cv

I. IS THE ZERO-POINT ENERGY OF QUANTIZED FIELDS OBSERVABLE?

General relativity theory (GRT), and the relativistic quantum field theories (QFT) of the standard model of elementary particles, are describing all experimental observations with impressive accuracy — as long as GRT or QFT are used separately. But as soon as one tries to combine these successful theories, serious problems turn up. One of the most spectacular examples of such incompatibility has been dubbed "the cosmological constant problem". A by now classical review article on that issue has been compiled by Weinberg [1]. See the article by Li *et. al.* [2] for an updated review. In short, the cosmological constant problem manifests itself as follows:

According to the field equation

$$R_{\mu\nu} - \frac{R}{2}\, g_{\mu\nu} + \Lambda\, g_{\mu\nu} = -\frac{8\pi G}{c^4}\, T_{\mu\nu} \tag{1}$$

of GRT, the curvature of space-time, represented by the Ricci-tensor ($R_{\mu\nu}$) and it's contraction R, is proportional to the energydensity-stress-tensor — henceforth named es-tensor for brevity — ($T_{\mu\nu}$), which again is determined by the energy density and the momentum density of all fields contained within space-time, i. e. of all fields with exception of the metric field ($g_{\mu\nu}$).

Upon canonical quantization of any classical continuous field, the Hamilton operator of that field will diverge. For example, canonical quantization of the classical electromagnetic field results into the Hamilton operator

$$H = \sum_{\boldsymbol{k}} \sum_{v=1}^{2} \frac{\hbar c\, |\boldsymbol{k}|}{2} \Big(a_{\boldsymbol{k}}^{(v)+} a_{\boldsymbol{k}}^{(v)} + a_{\boldsymbol{k}}^{(v)} a_{\boldsymbol{k}}^{(v)+} \Big) \tag{2a}$$

$$= \sum_{\boldsymbol{k}} \sum_{v=1}^{2} \hbar c\, |\boldsymbol{k}| \Big(a_{\boldsymbol{k}}^{(v)+} a_{\boldsymbol{k}}^{(v)} + \frac{1}{2} \underbrace{[a_{\boldsymbol{k}}^{(v)}, a_{\boldsymbol{k}}^{(v)+}]}_{1} \Big)\ , \tag{2b}$$

with $a_{\boldsymbol{k}}^{(v)+}$ and $a_{\boldsymbol{k}}^{(v)}$ being the creation- and annihilation-operators respectively of photons with wavenumber $\boldsymbol{k}$ and polarization v. The summation is running over all of the infinitely many wavenumbers $\boldsymbol{k}$, which are compatible with the normalization volume (if an infinite normalization volume is chosen, the sum is replaced by an integral). Due to the commutator, the energy is infinite. The waves described by the second term in (2b) are the zero-point-oscillations, and their energy is the zero-point-energy, of the quantized electromagnetic field.

The observed curvature of intergalactic space is close to zero [3], suggesting that either
(a) the cosmological constant Λ in (1) should be adjusted, to compensate the zero-point-energy of the quantum fields, or that
(b) the zero-point-energy of the quantum fields should be considered as "not existing", or that
(c) some combination of (a) and (b) should be considered.

Alternative (a) calls for a fine-tuning of the cosmological constant Λ with an accuracy of many dozens of decimal digits, the exact number depending on the method applied for regularization of the diverging term in (2b). Therefore this solution — though being completely correct under purely formal criteria — does seem to be quite "unnatural", and is not considered acceptable by many scientists.

On first sight, there seem to be less objections against alternative (b). In pure quantum-field-theoretical computations (neglecting gravity), only energy differences matter, but not absolute energy values. Therefore the offset of an infinitely large zero-point energy is merely a tiresome ballast without discernible functionality. To get rid of that offset, normal order is often applied as an ad-hoc measure. It means, that in (2a) all creation operators are shifted left, and all annihilation operators are shifted right, under disregard of their commutation relations! Thus the Hamilton-operator becomes

$$H = \sum_{\boldsymbol{k}} \sum_{v=1}^{2} \hbar c\, |\boldsymbol{k}|\, a_{\boldsymbol{k}}^{(v)+} a_{\boldsymbol{k}}^{(v)}\ , \tag{2c}$$

and the infinite energy offset has disappeared.

As the results of quantum field theoretical computations are not changed, if normal order is applied to the Hamilton operator, and as no gravitational effect of the zero-point-energy of quantum fields is observed, one might very well ask whether that zero-point-energy does

---

* e-mail: gerold.gruendler@astrophys-neunhof.de

exist at all. Words like "existence" or "reality" in this context of course mean the question, whether the zero-point-energy is observable and can be tested experimentally.

A possible method to observe the quantized electromagnetic field's zero-point energy — and actually the only method proposed until today, besides the missing gravitational effect — has been suggested by Casimir [4] in 1948. According to Casimir's computation (on which we will dwell below), the zero-point oscillations cause an attractive force

$$F_{\text{Casimir}} = -\frac{\pi^2 \hbar c}{240}\,\frac{XY}{D^4} = -1.3 \cdot 10^{-9}\text{N} \cdot \frac{XY/\text{mm}^2}{D^4/\mu\text{m}^4} \quad (3)$$

between two parallel metallic plates with area XY, which are separated by a gap of width D. Since then, the Casimir-force has many times been measured, and the — at least approximate — correctness of equation (3) has been confirmed [5].

These observations still do not conclusively prove the reality of the zero-point energy, because a completely different explanation for the same observations is available: Lifshitz [6] and Dzyaloshinskii, Lifshitz, and Pitaevskii [7] have computed the long-range van der Waals-forces, which are acting between two infinitely extended half-spaces with relative dielectric constants ϵ_1 and ϵ_3, which are separated by a gap filled with a material with relative dielectric constant ϵ_2. Schwinger, DeRaad, and Milton [8] reproduced and confirmed the results of Lifshitz *et. al.*. They also considered the limit $\epsilon_1 = \epsilon_3 \to \infty$, $\epsilon_2 \to 1$, i. e. the limit of two metal plates with infinite conductivity, separated by a vacuum gap. It turned out that the formula of Lifshitz *et. al.* simplifies in this limit to the Casimir-force (3).

Thus two different explanations exist for the same experimental results, and it seems to be merely a matter of taste, which one to prefer. Jaffe [9] called Casimir's method of derivation of equation (3) "heuristic", as the assumption of infinite conductivity of the metal plates (which is equivalent to taking the limit $\alpha \to \infty$, with α being the QED coupling constant) is obscuring the true nature of the interaction between the electromagnetic field and the charged matter in the metallic plates. While this criticism certainly is justified, even a heuristic argumentation may lead to correct results eventually.

In an attempt to clarify the validity and significance of Casimir's approach, in this article the important difference between the two methods of derivation of the force inbetween the metal plates is addressed. While Lifshitz *et. al.* did the complete computation consistently with finite parameters ϵ_1, ϵ_2, ϵ_3, Casimir started his computation with the assumption of finite conductivity of the metal plates, and changed only at the very end of the computation to the limit $\epsilon \to \infty$, to arrive at equation (3). Experimentalists then again must apply corrections to Casimir's formula, reflecting the final conductivity of the metals used in their experiments, to achieve good fits of their results [5].

What would be the outcome of Casimir's method of computation, if the detour via the assumption of infinite conductivity of the metals is avoided? The answer to this question, which is detailed in the following sections, is quite surprising: The experimental confirmation of Casimir's formula (3) is proving — exactly opposite to Casimir's interpretation — that the electromagnetic field's zero-point energy does *not* exert forces onto metallic plates, i. e. that the observed forces by no means are related to the electromagnetic field's zero-point energy.

II. CASIMIR'S COMPUTATION



FIG. 1. Cavity resonator with movable plate

Casimir considered a resonator as sketched in figure 1. The rectangular cavity's size is $X \times Y \times (Z + P)$. Inside the cavity there is a plate of thickness P, which is aligned parallel to the cavity's XY-face and movable in Z-direction. The plate's distance from one side wall of the cavity is D, it's distance from the opposite side wall is $Z - D$.

We firstly consider the left cavity. At temperature T, it's walls and the plate are in thermodynamic equilibrium with the electromagnetic blackbody radiation within the cavity. The spectrum of the radiation's wave numbers is discrete:

$$k_{rst} = \sqrt{\left(\frac{r\pi}{X}\right)^2 + \left(\frac{s\pi}{Y}\right)^2 + \left(\frac{t\pi}{D}\right)^2}$$
$$\text{with } r, s, t \in \mathbb{N} \quad (4)$$

There are 2 modes each with $r, s, t = 1, 2, 3, \ldots$ and 1 mode each with one of the indices 0 and the both other indices $1, 2, 3, \ldots$ [10, chap. D.II.2.b.].

This equation does not hold for arbitrary wave numbers, because any metal becomes transparent for radiation of sufficiently high frequency. The transparency of metals at very high frequency is a basic condition for Casimir's derivation of formula (3). Casimir was well aware of this condition, which he emphasized in his publication [4]. He introduced "a function $f(k_{rst}/k_{\text{M}})$, which is unity for $k_{rst} \ll k_{\text{M}}$ but tends to zero sufficiently rapidly for

$k_{rst} \to \infty$". But he did not specify the concrete form of $f(k_{rst}/k_{\mathrm{M}})$ explicitly.

To design $f(k_{rst}/k_{\mathrm{M}})$ realistic, we assume the cavity and the plate to be made of copper. The reflectivity R as a function of wave number is quite complicated, see [11, fig. 1b]. For the purpose of our investigation, the rough approximation

$$R = \exp\{-\,k_{rst}/k_{\mathrm{M}}\} \quad \text{with } k_{\mathrm{M}} = 38 \cdot 10^6 \mathrm{m}^{-1} \tag{5}$$

for the reflectivity of copper is completely sufficient. (k_{M} is about 0.8× the plasma-wavenumber of copper [11] according to the Drude-model.)

At temperature $T = 0$, only the zero-point oscillations of the electromagnetic field are excited. The energy per mode then is $\hbar c k_{rst}/2$, and the zero-point energy enclosed within the cavity is

$$U_0 = 2 \sum_{r,s,t=0}^{\infty}{}' \, \frac{\hbar c k_{rst}}{2} \exp\{-k_{rst}/k_{\mathrm{M}}\} \; . \tag{6}$$

The prime$'$ at the summation symbol is a reminder, that the multiplicity of polarizations, as indicated in (4), has to be considered. At most one of the numbers r, s, t of an oscillation mode can be zero, and terms with one zero index get a factor 1/2.

Casimir considered the limit $X \to \infty$ and $Y \to \infty$. Thus he could replace the sums over the discrete indices r and s by integrals. Only the sum over t is still considered discrete. It has been elaborated elsewhere [12, sect. 4] in very detail, how (6) then can be transformed into

$$\begin{aligned} U_0 = \frac{\pi^2 \hbar c X Y}{2} \bigg(& \frac{6 D k_{\mathrm{M}}^4}{\pi^4} - \frac{1}{360\, D^3} + \\ & + \sum_{j=6}^{\infty} \frac{B_j}{j!} \, \frac{(j^2 - 5j + 6)\, \pi^{j-4}}{k_{\mathrm{M}}^{j-4} D^{j-1}} \bigg) \; . \end{aligned} \tag{7}$$

The coefficients B_j are the Bernoulli-numbers

$$\begin{aligned} B_0 &= 1 \\ B_j &= -\sum_{n=0}^{j-1} \frac{j!}{n!(j-n+1)!}\, B_n \quad \text{for } j > 0 \; . \end{aligned} \tag{8}$$

The expansion, which led to the series with the Bernoulli coefficients, does converge only for

$$D > \frac{1}{2 k_{\mathrm{M}}} \overset{(5)}{=} 13.2\,\mathrm{nm} \; . \tag{9}$$

For smaller distance D between the movable plate and the cavity wall, (7) is not valid.

(7) is the zero-point energy enclosed within the left cavity of figure 1. The energy within the right cavity is found due to replacing D everywhere by $Z - D$. Then the force

$$F_{\mathrm{ZPE}} = -\frac{\mathrm{d}U_{0,\mathrm{total}}}{\mathrm{d}D} \tag{10}$$

acting onto the movable plate, which is caused by the different content of zero-point energy in the both cavities, can be computed. (The index $_{\mathrm{ZPE}}$ stands for zero-point energy.) With the approximation

$$\left(\frac{D}{Z - D} \right)^4 \ll 1 \; , \tag{11}$$

which is well justified for all experimental evaluations of the Casimir force, the result is

$$\begin{aligned} F_{\mathrm{ZPE}} = -\frac{\pi^2 \hbar c X Y}{2} \bigg(& \frac{6 (k_{\mathrm{M,left}}^4 - k_{\mathrm{M,right}}^4)}{\pi^4} + \frac{1}{120\, D^4} + \\ & + \underbrace{\sum_{j=6}^{\infty} \frac{B_j}{j!} \, \frac{(-j^3 + 6j^2 - 11j + 6)}{k_{\mathrm{M}}^{j-4}\, \pi^{4-j} D^j}}_{A} \bigg) \; . \end{aligned} \tag{12}$$

Here different values $k_{\mathrm{M,left}}$ and $k_{\mathrm{M,right}}$ have been inserted, considering the unavoidable differences in reflectivity due to surface contaminations, scratches, and the like. This will be discussed in detail in the next section. Casimir did *not* consider such possible differences. Instead at this point of the computation, he switched to the limit $k_{\mathrm{M,left}} = k_{\mathrm{M,right}} \to \infty$, which immediately led to his formula (3).

Skipping the last term, named A in (12), indeed is justified in most cases, because the Bernoulli-expansion is converging rapidly:

$$\frac{A}{1/(120\, D^4)} \lesssim \begin{cases} 5\,\% & \text{for } D > 0.2\,\mu\mathrm{m} \\ 1\,\% & \text{for } D > 0.5\,\mu\mathrm{m} \end{cases} \tag{13}$$

But it was a misleading idealization, to neglect the k_{M}^4 term.

III. KEEPING PARAMETERS REALISTIC

Due to the idealizing assumption $k_{\mathrm{M,left}} = k_{\mathrm{M,right}}$, Casimir dropped the term $6(k_{\mathrm{M,left}}^4 - k_{\mathrm{M,right}}^4)/\pi^4$. This term is describing the fragile balance of two forces, which are tremendous in comparison to the Casimir-force $\sim 1/(120\, D^4)$. Using the realistic parameter $k_{\mathrm{M}} \overset{(5)}{=} 38 \cdot 10^6 \mathrm{m}^{-1}$, the ratio of these forces is

$$\frac{6 k_{\mathrm{M}}^4/\pi^4}{1/(120\, D^4)} \approx \begin{cases} 2.5 \cdot 10^4 & \text{for } D = 0.2\,\mu\mathrm{m} \\ 1.5 \cdot 10^7 & \text{for } D = 1.0\,\mu\mathrm{m} \\ 9.6 \cdot 10^9 & \text{for } D = 5.0\,\mu\mathrm{m} \end{cases} \tag{14}$$

Only with excellent, and actually not realistic match of reflectivities of the both sides of the plate, the Casimir force can become visible. For $D = 1\,\mu$m, the ratio of the forces is

$$\begin{array}{ccc} \dfrac{k_{\mathrm{M,right}}}{k_{\mathrm{M,left}}} & & \dfrac{F_{\mathrm{Casimir}} = (3)}{F_{\mathrm{ZPE}} = (12)} \\ 1 - 10^{-4} & \Longrightarrow & 1.6 \cdot 10^{-4} \\ 1 - 10^{-5} & \Longrightarrow & 1.6 \cdot 10^{-3} \\ 1 - 10^{-6} & \Longrightarrow & 1.6 \cdot 10^{-2} \\ 1 - 10^{-7} & \Longrightarrow & 1.4 \cdot 10^{-1} \\ 1 - 10^{-8} & \Longrightarrow & 6.2 \cdot 10^{-1} \end{array} \tag{15}$$

Actually, the experimentalists don't even try to achieve good matching surface conditions on the both sides of the movable plate. For example in the Purdue experiment [13, 14], the wall of the cavity is replaced by an Au-coated sphere, and the movable plate is replaced by a 3 μm thick silicon plate, which is Cu- or Au-coated on it's side facing the sphere, but un-coated bare Si on it's rear side. In a Yale experiment [15], the cavity wall again is replaced by an Au-coated sphere, while the movable plate is replaced by a SiN membrane, which is Au-coated on it's side facing the sphere, but un-coated bare SiN on it's rear side.

Still these experiments — like all other experiments, which as well don't pay attention to the surface conditions of their movable plate's rear sides — are seeing the Casimir-force (3), but not the force $F_{\mathrm{ZPE}} = (12)$ caused by zero-point energy.

If the zero-point energy would exert forces onto metallic surfaces, then the term $k^4_{\mathrm{M,left}} - k^4_{\mathrm{M,right}}$ would dominate the observed forces in all of these experiments, making the tiny $\sim D^{-4}$ Casimir-force invisible. F_{ZPE} does not depend on D, and it can — depending on the relative reflectivities of the two sides of the movable plate — be attractive or repulsive. No force with that signature has ever been reported from any Casimir-force experiment.

Thus the experimental observations of the Casimir-force (3) do stringently prove, that the force $F_{\mathrm{ZPE}} = (12)$ (caused by the electromagnetic field's zero-point oscillations) does not exist.

This result saves us from an intricate consideration: If a real photon impinges onto a metal surface, it may be reflected, or it may be absorbed, or it may be transmitted. In contrast, zero-point energy obviously cannot be absorbed. Otherwise any piece of matter, which is surrounded by the electromagnetic field's ubiquitous zero-point oscillations, would absorb by and by short-wavelength zero-point radiation, convert it into high-energy phonons, which within soon would thermalize into many low-energy phonons. Eventually, the matter would emit part of that energy again in form of low-energy *real* photons. Thus matter would permanently convert high-energy zero-point photons into lower-energy real photons, thereby heating up to infinity.

In his final result, Casimir switched to infinite conductivity of the metal, thus excluding absorption. But in course of his derivation, as he clearly pointed out in his publication [4], Casimir assumed that most of the long-wavelength zero-point oscillations are reflected by the cavity walls, and most of the very-short-wavelength zero-point oscillations are transmitted. But he never assumed that absorption might happen at any wavelength. This is a very strange, and almost contradictory assumption, because — if considered on a microscopic scale — the processes of absorption and reflection of electromagnetic radiation are almost identical. Therefore it seems to be a far-fetched idea, that a real metal, which is able to reflect some zero-point electromagnetic radiation, should not be able to absorb that same zero-point radiation.

IV. CONCLUSIONS

The findings presented in this article certainly do not solve the cosmological constant problem. But they may be helpful to avoid blind alleys, and to stir the search for a solution into the right direction. The essential facts are: Firstly, no gravitational effect, caused by the zero-point energy of quantum fields, has been observed. Secondly, the results of QFT — including all of the impressive achievements of QED like Lamb-shift, electron g-factor, hydrogen hyperfine-splitting, and so on — can be found, if normal order (2c) is applied to the Hamilton operator. Thirdly, as shown in this article, the observed Casimir-force is definitely not caused by zero-point energy.

In total, no experimental evidence at all is indicating the measurable, observable existence of the zero-point energy of quantum fields. Therefore, instead of renormalizing the cosmological constant Λ (or even modifying GRT), it is certainly more promising, to approach the problem directly at it's root, i. e. to somehow remove zero-point energy from QFT. The crude measure of normal order is not an acceptable solution, as the disregard of the non-commutative operator algebra is irreconcilable with the basic principles of QFT. Unfortunately, at this moment I cannot offer a better solution.

ACKNOWLEDGMENTS

This work gained much from innumerable helpful discussions on the mysterious quantum vacuum with my colleagues V. I. Nachtmann and O. S. ter Haas.

[1] S. Weinberg, Rev. Mod. Phys., **61**, 1 (1989).
[2] M. Li, X.-D. Li, S. Wang, and Y. Wang, Commun. Theor. Phys., **56**, 525 (2011).
[3] C. L. Bennet *et. al.*, arXiv [astro-ph.CO]: 1212.5225 (2012).
[4] H. B. G. Casimir, Proc. Kon. Nederl. Akad. Wetensch., **51**, 793 (1948),

http://www.astrophys-neunhof.de/serv/Casimir1948.pdf.

[5] A. Lambrecht and S. Reynaud, Int. J. Mod. Phys. Conf. Ser., **14**, 171 (2012).

[6] E. M. Lifshitz, Sov. Phys. JETP, **2**, 73 (1956).

[7] I. E. Dzyaloshinskii, E. M. Lifshitz, and L. P. Pitaevskii, Sov. Phys. Usp., **73**, 381 (1961).

[8] J. Schwinger, L. L. DeRaad, and K. A. Milton, Ann. Phys. (NY), **115**, 1 (1978).

[9] R. L. Jaffe, Phys. Rev. D, **72**, 021301 (2005).

[10] F. W. Gundlach, *Grundlagen der Höchstfrequenztechnik* (Springer, Berlin, 1959).

[11] H. Ehrenreich and H. R. Philipp, Phys. Rev., **128**, 1622 (1962).

[12] G. Gründler, APIN Circular se08011 (2013), http://www.astrophys-neunhof.de/mtlg/se08011.pdf.

[13] R. S. Decca, D. López, E. Fischbach, and D. E. Krause, Phys. Rev. Lett., **91**, 050402 (2003).

[14] R. S. Decca, D. López, E. Fischbach, G. L. Klimchitskaya, D. E. Krause, and V. M. Mostepanenko, Phys. Rev. D, **75**, 077101 (2007).

[15] D. Garcia-Sanchez, K. Y. Fong, H. Bhaskaran, S. Lamoreaux, and H. X. Tang, Phys. Rev. Lett., **109**, 027202 (2012).